



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The presence and role of hypoxia in the endometrium

Citation for published version:

Martínez-aguilar, R, Kershaw, LE, Reavey, JJ, Critchley, HOD & Maybin, JA 2021, 'The presence and role of hypoxia in the endometrium', *Reproduction*, vol. 161, no. 1. <https://doi.org/10.1530/REP-20-0268>

Digital Object Identifier (DOI):

[10.1530/REP-20-0268](https://doi.org/10.1530/REP-20-0268)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Reproduction

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



The presence and role of hypoxia in the endometrium

Rocio Martínez-Aguilar¹, Lucy E. Kershaw^{2,3}, Jane J. Reavey¹, Hilary O.D. Critchley¹ and Jacqueline A. Maybin^{1*}.

¹MRC Centre for Reproductive Health, University of Edinburgh, The Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh, UK

²Edinburgh Imaging, University of Edinburgh, The Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh, UK

³Centre for Inflammation Research, University of Edinburgh, The Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh, UK

* Corresponding author:

Dr Jacqueline A Maybin

MRC Centre for Reproductive Health

The University of Edinburgh

The Queen's Medical Research Institute

47 Little France Crescent

EDINBURGH

UK

Email: Jackie.maybin@ed.ac.uk

Short Title: Hypoxia in the endometrium

Word Count: 7552

ABSTRACT

The endometrium is a multicellular tissue that is exquisitely responsive to the ovarian hormones. The local mechanisms of endometrial regulation to ensure optimal function are less well characterised. Transient physiological hypoxia has been proposed as a critical regulator of endometrial function. Herein, we review the literature on hypoxia in the non-pregnant endometrium. We discuss the pros and cons of animal models, human laboratory studies and novel *in vivo* imaging for the study of endometrial hypoxia. These research tools provide mounting evidence of a transient hypoxic episode in the menstrual endometrium and suggest that endometrial hypoxia may be present at the time of implantation. This local hypoxia may modify the inflammatory environment, influence vascular remodelling and modulate endometrial proliferation to optimise endometrial function. Finally, we review current knowledge of the impact of this hypoxia on endometrial pathologies, with a focus on abnormal uterine bleeding. Throughout the manuscript areas for future research are highlighted with the aim of concentrating research efforts to maximise future benefits for women and society.

39	Table of Contents	
40	Abstract	3
41	Introduction	5
42	Endometrial histology and ovarian hormone regulation.....	5
43	Endometrial breakdown and regeneration	6
44	Detection of Hypoxia throughout the menstrual cycle	7
45	In vivo animal models	7
46	Rodent models	7
47	Macaque models.....	12
48	Ex vivo human endometrial studies.....	13
49	In vivo human endometrial studies	13
50	Role of hypoxia throughout the menstrual cycle.....	15
51	Impact of hypoxia on inflammation.....	16
52	Impact on neutrophils.....	16
53	Impact on macrophages.....	18
54	Impact of hypoxia on proliferation	20
55	Impact of hypoxia on vascular remodelling and angiogenesis.....	21
56	The role of hypoxia in endometrial pathology	23
57	Abnormal uterine bleeding.....	23
58	Structural causes of AUB.....	23
59	Non-structural causes of AUB	27
60	Conclusions	28
61	References	30
62	FIGURE LEGENDS.....	44
63		
64		

INTRODUCTION

The human endometrium is a heterogeneous and dynamic tissue that undergoes cyclical breakdown and repair/regeneration more than 400 times during the female reproductive lifespan (Short, 1976; Critchley *et al.*, 2020). This occurs each month without scarring or loss of function. However, the regulation and local mechanisms of this endometrial breakdown and repair remain elusive. In particular, our knowledge of the contribution of local endometrial hypoxia to this process is in its infancy. The presence of hypoxia, usually defined as a partial oxygen pressure below 10 mmHg, is not an uncommon phenomenon in human physiology, e.g. bone marrow and intestinal mucosa (Suda, Takubo & Semenza, 2011; Zheng, Kelly & Colgan, 2015). Its presence in the menstrual endometrium has been proposed following progesterone withdrawal and intense vasoconstriction of the specialised spiral arterioles (Markee, 1940). Unravelling the role of hypoxia in the endometrium has the potential to improve our understanding of menstrual and implantation disorders and reveal novel therapeutic strategies for those suffering from these common, devastating conditions.

ENDOMETRIAL HISTOLOGY AND OVARIAN HORMONE

REGULATION

Histologically, the endometrium can be divided into the functional and basal layer (Noyes, Hertig & Rock, 1950). The functional layer occupies the upper two thirds of the endometrium and is composed of stroma and glands. This layer undergoes constant remodelling throughout the menstrual cycle and is shed during menstruation. The basal layer, adjacent to the myometrium, comprises the lower third of the endometrium.

Oestradiol is the dominant hormone in the first half of the menstrual cycle, during the proliferative phase. It acts via the oestrogen receptor (ER), which has two structurally related subtypes, ER α and ER β (Lessey *et al.*, 1988; Critchley *et al.*, 2002). After ovulation, levels of oestradiol decline and the corpus luteum increases its progesterone production, prompting endometrial differentiation and decidualisation. This process, driven by cAMP signalling, reshapes the stromal compartment in order to keep the endometrium receptive for future implantation (Dunn, Kelly & Critchley, 2003). In contrast with non-menstruating species, where implantation of an embryo is required to trigger decidualisation (Brasted *et al.*, 2003), the human endometrium spontaneously decidualises with endometrial stromal cells in close proximity to spiral arterioles initiating their own transformation (Gellersen & Brosens, 2014). They morphologically transition from fibroblast-like cells to rounded epithelioid-like cells (Dunn, Kelly & Critchley, 2003).

ENDOMETRIAL BREAKDOWN AND REGENERATION

In the absence of implantation, the corpus luteum regresses causing significant progesterone withdrawal (Corker *et al.*, 1976; Maybin, Hirani, *et al.*, 2011). This decrease in progesterone levels triggers a cascade of local physiological inflammatory events that initiate menstruation. Progesterone withdrawal leads to the induction of the transcription factor NF κ B, which up-regulates the expression of pro-inflammatory cytokines (IL-6, TNF) and chemokines (CCL2, CXCL8)(King, Critchley & Kelly, 2001). In addition, this fall in progesterone levels increases endometrial cyclooxygenase 2 (COX-2), responsible for the synthesis of prostaglandins (PG)(Critchley *et al.*, 1999). Increased levels of these inflammatory mediators drive the recruitment of myeloid leukocytes, activation of matrix metalloproteinases (MMPs) and the shedding of the upper endometrial layers (Critchley *et al.*, 2001; Kelly, King & Critchley, 2001). Hypoxia has been identified in the endometrium

following progesterone withdrawal (Fan *et al.*, 2008; Cousins, Murray, *et al.*, 2016; Maybin *et al.*, 2018) and may be due to vasoconstriction of the endometrial vessels. $\text{PGF}_{2\alpha}$ and endothelin-1 (ET-1) are two endometrial factors with known vasoconstrictive properties that are present following progesterone withdrawal (Baird *et al.*, 1996; Marsh *et al.*, 1997). Vasoconstriction of specialised endometrial spiral arterioles may limit blood loss during menstruation. The subsequent tissue hypoxia does not appear to be necessary for endometrial breakdown but may have an important role in endometrial repair/regeneration (Maybin *et al.*, 2018; Chen *et al.*, 2020).

Shedding of the functional endometrial layer necessitates repair of the denuded endometrial surface and regeneration of endometrial tissue. This takes place when oestradiol and progesterone levels are low but local glucocorticoid action may be increased (McDonald *et al.*, 2006; Kaitu'u-Lino, Morison & Salamonsen, 2007a; Rae *et al.*, 2009). Evidence from mouse models and human tissue studies suggest that hypoxia is required for physiological endometrial repair (Fan *et al.*, 2008; Maybin *et al.*, 2018). The processes involved are likely to be similar to those of wound healing, involving haemostasis, inflammation, proliferation and remodelling (Velnar, Bailey & Smrkolj, 2009; Mutsaers *et al.*, 2015).

DETECTION OF HYPOXIA THROUGHOUT THE MENSTRUAL CYCLE

The first suggestion that hypoxia was present at menses derived from findings in a primate model in 1940 (Markee, 1940). Transplantation of endometrial explants to the Rhesus macaque eye allowed direct observation of intense vasoconstriction of spiral arterioles and focal bleeding following

progesterone withdrawal. Since then, the use and refinement of animal models for the study of menstrual physiology and endometrial hypoxia has become more common.

In vivo animal models

Menstruation is restricted to humans and few other species. These include higher order primates (baboons, Rhesus macaques), the elephant shrew (Van der Horst & Gillman, 1940), certain bats (Hamlett, 1934; Rasweiler & de Bonilla, 1992; Zhang *et al.*, 2007) and the spiny mouse (Bellofiore *et al.*, 2017). The majority of menstrual studies have been carried out in rodents and non-human primates, including the Rhesus macaque (Brenner & Slayden, 2012).

Rodent models

Despite physiological differences between mice and humans (e.g. a shorter length of cycle and lack of spontaneous decidualisation) mouse models replicate the events of human menstruation and decidualisation well (Wang *et al.*, 2013; Cousins, Kirkwood, *et al.*, 2016; Armstrong *et al.*, 2017). The feasible management of large experimental groups, short breeding times and availability of laboratory antibodies/reagents provide advantages over macaque models. Mouse models also offer the possibility of genetic, environmental and pharmacological manipulation of hypoxia (see [Role of hypoxia throughout the menstrual cycle](#) below). Technically, euthanasia by carbon dioxide (CO₂) inhalation can impact tissue hypoxia and may distort results. Hence, cervical dislocation is the recommended euthanasia method for these studies. Great care must be taken to handle, process and fix tissue rapidly to capture the physiological events of menstruation.

The mouse model of simulated menstruation

The *menses-like* model was first described in 1984 (Finn & Pope, 1984) and further optimised in the 2000's (Brasted *et al.*, 2003). Since then, it has been the most popular model to investigate the dynamics of endometrial repair (Fan *et al.*, 2008; Evans, Kaitu'u-Lino & Salamonsen, 2011; Cousins *et al.*, 2014; Maybin *et al.*, 2018; Chen *et al.*, 2020) (**Fig. 1**). Mice are ovariectomised and supplemented with exogenous oestradiol and progesterone to mimic the human hormonal endometrial environment. They require artificial induction of decidualisation, via a transcervical or surgical intrauterine injection of oil. Once decidualisation has taken place, progesterone withdrawal leads to active bleeding in the mouse uterus and subsequent repair (**Fig. 1a**). Alternatively, simulation of menses can be achieved by inducing pseudopregnancy (**Fig. 1b**). In this model, female mice are mated with vasectomized males to mimic fertilisation events. Progesterone withdrawal occurs naturally or is induced by ovariectomy or administration of a progesterone antagonist (Rudolph *et al.*, 2012).

The first work to describe the presence of hypoxia during endometrial breakdown and repair in the mouse utilised the 'pseudopregnancy' model variant (Fan *et al.*, 2008). Pimonidazole is a hypoxic marker that, when oxygen partial pressures are below 10 mmHg, forms protein adducts which can be visualized using specific monoclonal antibodies. Due to its chemical stability, pimonidazole is considered one of the most reliable means of tissue oxygen level detection, even when it is temporally and spatially transient. Fan *et al.* found the endometrial area undergoing regeneration to be hypoxic and that this hypoxia decreased and eventually disappeared with endometrial reepithelialisation (Fan *et al.*, 2008). Subsequent confirmation of the presence of menstrual hypoxia was found in the 'exogenous hormone' model of simulated menses (Cousins, Murray, *et al.*, 2016; Maybin *et al.*, 2018; Chen *et al.*, 2020). Using pimonidazole, hypoxia was detected during bleeding and later confined to areas undergoing active repair. Hypoxia may also be present in the endometrium at the time of implantation. As the uterine epithelium contains no blood vessels

during initial embryo contact, it has been suggested that the onset of implantation occurs in a hypoxic environment (Daikoku *et al.*, 2003). The detection of pimonidazole adducts in the area of implantation in mice reinforces this hypothesis (Pringle *et al.*, 2007).

Another method to determine tissue hypoxia is detection of the oxygen-sensing transcription factor hypoxia inducible factor (HIF). HIFs have a key role in the cellular response to oxygen and are heterodimers composed of two subunits: a constitutively expressed beta subunit (HIF-1 β) and an O₂-sensitive alpha subunit (Semenza, 2000). There are three known α subunits: HIF-1 α , HIF-2 α , and HIF-3 α . HIF-1 α and HIF-2 α are the most common alpha isoforms and present overlapping but distinct target gene specificities (Mole *et al.*, 2009). HIF-3 α is structurally different from the other isoforms and is the least characterized (Pasanen *et al.*, 2010). Along with promoting genes related to nitrogen metabolism and immune response, HIF-3 α has the ability to inhibit HIF-1 α /2 α action (Zhang *et al.*, 2014).

The regulation of HIF takes place predominantly at the protein level. In normoxia, prolyl hydroxylase domain enzymes (PHDs) hydroxylate specific residues within the alpha subunit, leading to its ubiquitination and subsequent degradation via the proteasome (Salceda & Caro, 1997). In hypoxia these PHDs are inhibited, resulting in HIF- α stabilization. HIF- α translocates to the nucleus, dimerizes with HIF-1 β and binds to hypoxia-response elements (HREs) to enhance transcription of a plethora of genes involved in energy metabolism, angiogenesis, tissue remodelling and inflammatory responses (Semenza, 2012).

The presence of nuclear HIF-1 α protein is therefore indicative of active HIF-1 and consistent with tissue hypoxia. Using this approach, HIF-1 α has been detected during menstruation in both the exogenous hormone (Maybin *et al.*, 2018; Chen *et al.*, 2020) and pseudopregnancy menstruation models (Chen *et al.*, 2015), decreasing during endometrial regeneration. Examination of HIF-1 α and

HIF-2 α in the mouse uterus during pre-implantation (day 4) and decidualisation (day 5-8) of pregnancy, revealed HIF-1 α was present in the luminal epithelium prior to implantation and throughout the epithelium and stroma during decidualisation and implantation (Daikoku *et al.*, 2003). HIF-2 α was seen in the stroma on day 4 and limited to cells surrounding the blastocyst on day 5. The authors suggested that HIF-1 was involved in maintaining oxygen homeostasis and that HIF-2 was driving the angiogenesis necessary for successful implantation.

Various concerns have been raised about using HIF as a hypoxic surrogate marker. Transient hypoxic events can be too brief to stabilise HIF for immunohistochemical detection (Wang *et al.*, 1995). Antibody unreliability is an added factor, which is compounded by the fact that tissue collection and fixation can also affect HIF detection (Zhang & Salamsen, 2002). Furthermore, HIF stabilisation can be induced by NF- κ B-driven cytokine production in a non-hypoxic dependent manner and hypoxia can exert downstream effects independently of HIF signalling (Lin & Simon, 2016). Alongside detection of pimonidazole and HIF, hypoxia-inducible factor downstream targets may indicate a hypoxic response in the mouse menstrual endometrium. HIF-1 α -mediated induction of the angiogenic factors vascular endothelial growth factor (VEGF) and the chemokine receptor CXCR4 was increased during menstruation and endometrial repair (Fan *et al.*, 2008; Chen *et al.*, 2015; Cousins, Murray, *et al.*, 2016; Maybin *et al.*, 2018).

Xenograft mouse model

The xenograft mouse model provides an alternative model for study of menstrual physiology and pathology (extensively reviewed in (Kuokkanen, Zhu & Pollard, 2017)). Human functional endometrium is transplanted into immunodeficient mice (**Fig. 1c**). This is usually collected during the proliferative phase and can be transplanted as (i) small fragments (1-2 mm³) of endometrial tissue

(Guo *et al.*, 2011; Coudyzer *et al.*, 2013) or (ii) dissociated endometrial cells from epithelial and stromal fractions that are mixed before implantation (Masuda *et al.*, 2007; Polotsky *et al.*, 2009). The recipient mice are selected to limit xenograft tissue rejection, but the immunodeficient strain used can vary. The most commonly used in xenograft menstruation models is the severe combined immunodeficiency (SCID) mouse, which has T and B cell deficiencies (Gaide Chevronnay *et al.*, 2009; Guo *et al.*, 2011; Coudyzer *et al.*, 2013). The best engraftment results are achieved with the non-obese diabetic (NOD)/SCID/ γc^{null} mice (NOG), which also have defective NK cell activity (Matsuura-Sawada *et al.*, 2005; Masuda *et al.*, 2007).

Generally, the patches of endometrial tissue are placed subcutaneously in mice (Guo *et al.*, 2011; Coudyzer *et al.*, 2013) with a survival time of 4 weeks, whereas the dissociated endometrial cells are implanted below the kidney capsule and survive up to 10 weeks (Masuda *et al.*, 2007). This latter mode of implantation allows extension of the duration of experiments, making this the method of choice for studies of the proliferation kinetics of the endometrium after pharmacological treatments (Polotsky *et al.*, 2009).

Xenograft menstruation studies mainly focus on endometrial regeneration and the role of ovarian steroids in orchestrating the process (Gaide Chevronnay *et al.*, 2009; Guo *et al.*, 2011; Coudyzer *et al.*, 2015) and use the endometrial fragments model variant. To date, this mouse model has only been employed once to study the presence of hypoxia during menstruation (Coudyzer *et al.*, 2013). In 2013, Coudyzer *et al.* subcutaneously implanted endometrial patches on SCID female mice and tested for signs of hypoxia in the resulting xenograft using several methods. Firstly, they directly measured the local partial oxygen pressure (pO_2) using electron paramagnetic resonance and OxyLite fluorescent probes. They also studied the presence of pimonidazole staining and HIF-1 α using immunohistochemistry (IHC). The authors did not detect hypoxia during endometrial breakdown or repair using any of these methods. These results contrast with findings in the mouse

model of simulated menses and may be partially explained by the xenograft model itself. Endometrial tissue architecture and vasculature is severely compromised following transplantation and may impair vasoconstriction and prevent endometrial hypoxia. Moreover, endometrial breakdown and repair are considered inflammatory events, as they involve pro-inflammatory cytokine production and myeloid leukocyte recruitment (Finn, 1986). Therefore, the necessary immunosuppressed state of the recipient mice may alter physiological menstrual endometrial events. The SCID model aims to suppress T and B-cell mediated transplant/xenograft rejection without substantially affecting the innate immune response and may be more relevant than other immunocompromised recipient mice (Guo *et al.*, 2011; Donoghue *et al.*, 2012).

Spiny mouse

The common spiny mouse (*Acomys cahirinus*) is, to date, the only known rodent to display spontaneous decidualisation and natural menstruation (Bellofiore *et al.*, 2017, 2018). Although anatomically different, the spiny mouse uterus has physiological similarities to the human endometrium. For example, the spiny mouse displays spiral arteriole remodelling in the perimenstrual phase (Bellofiore *et al.*, 2018). In addition, endometrial decidualisation is tightly controlled, not compromising the structural integrity of the endometrial glands or the myometrium, as observed in other mouse models (Bellofiore *et al.*, 2018). Hypoxia has not been examined in this rodent to date and these studies are awaited with interest.

Macaque models

Macaques have morphologically similar uteri to humans, a similar length of menstrual cycle and they display spontaneous decidualisation (Brenner & Slayden, 2012). Macaques also experience menstrual abnormalities (e.g. heavy menstrual bleeding (HMB)) and can be fitted with tampons, hence they are exceptional candidates for evaluating therapies for menstrual disorders (reviewed in

(Brenner & Slayden, 2012)). Despite menstruating naturally, macaques are routinely ovariectomized and treated with oestradiol and progesterone to create artificial menstrual cycles and enable accurate timing of endometrial sampling. However, the need for larger experimental groups, longer experimental times and the increased cost of these experiments has meant many researchers are now preferentially using rodent models to study menstrual physiology.

As previously mentioned, the first indication of endometrial tissue hypoxia was observed in endometrial explants transplanted to the eye of rhesus macaques in the 1940s (Markee, 1940). Rather than hypoxia, Markee observed pulses of intense vasoconstriction in the spiral arterioles that he associated with localised hypoxic ischemia. This hypothesis was later supported by the detection and increased expression of HIF-1 α in the functional layer of the macaque endometrium during menstruation (Brenner & Slayden, 2012), consistent with the presence of endometrial hypoxia.

Ex vivo human endometrial studies

HIF-1 α protein has been identified, both by western blot and IHC, in human endometrial biopsies collected during the late secretory and menstrual phases (Critchley *et al.*, 2006; Maybin *et al.*, 2018). HIF-1 α staining was localised in the glandular and stromal cells in the functional endometrium, whereas in the basal layer HIF-1 α staining was restricted to the glands. In contrast, HIF-2 α is present exclusively during the early-mid secretory phase (Maybin *et al.*, 2018). Downstream targets of HIF, such as VEGF and carbonic anhydrase IX (CA-IX) have also been shown to be increased during the menstrual and proliferative phases (Stephen Charnock-Jones *et al.*, 1993; Sharkey *et al.*, 2000; Punyadeera, 2006; Maybin, Hirani, *et al.*, 2011).

In vivo human endometrial studies

Detection of human endometrial hypoxia *in vivo* has been largely via measurements of perfusion, initially investigated using thermal heat dissipation (Prill & Götz, 1961) and later by a Xenon-133 clearance technique (Fraser *et al.*, 1987) (**Fig. 2**). Both methods are invasive and results were conflicting as suffering from variable calibration and poor spatial and temporal resolution respectively. The introduction of Doppler ultrasound allowed perfusion measurements in individual spiral arterioles (Kupesic & Kurjak, 1993), but this showed an increase in flow the day before ovulation, in contrast with the ^{133}Xe clearance study which found a fall at this time. Laser Doppler fluxmetry was able to assess endometrial perfusion using a fibre optic probe (Gannon, Carati & Verco, 1997), finding blood flow peaks in the early proliferative and early secretory phase, but spatial resolution was limited. The more sensitive three-dimensional power Doppler angiography (3D-PDA) was also used in spiral arterioles (Raine-Fenning, 2004) and revealed a significant pre-ovulatory peak in perfusion, followed by a post-ovulatory fall and gradual increase through early to mid-secretory phases. In general, there has been little consensus regarding changes in endometrial blood flow over the menstrual cycle and how to measure such changes. Magnetic resonance imaging (MRI) methods may now offer a better alternative, although there has been little work on the application of these techniques to detect endometrial hypoxia.

To our knowledge, functional investigation of the normal endometrium has been limited to MR spectroscopy (Sarac *et al.*, 2004; Celik *et al.*, 2005). This technique detects the presence of specific metabolites in the body by examining the resonant frequencies of the hydrogen protons within them. In particular, lactate is a product of anaerobic respiration (and therefore a marker of hypoxia) and has been detected in normal secretory and proliferative endometrium (Sarac *et al.*, 2004; Celik *et al.*, 2005). Although lactate is arguably a more direct marker of hypoxia than measurement of

perfusion, analysis and acquisition of spectroscopy data is technically challenging (Lange *et al.*, 2006) and spatial resolution tends to be poor.

Dynamic contrast-enhanced (DCE) MRI is a technique that can detect hypoxia indirectly by measuring perfusion using an exogenous gadolinium-based contrast agent (CA) (Sourbron, 2010). Passage of the CA through the tissue can be modelled to allow perfusion to be estimated as part of a model-fitting process (Sourbron & Buckley, 2012). The technique has been applied in the normal endometrium (Majd *et al.*, 2017) but showed no differences between the secretory and proliferative phases. The advantage of DCE-MRI for hypoxia imaging is its good spatial resolution, but imaging and analysis can be complex (Brix *et al.*, 2004, 2009; Michaely *et al.*, 2008) and there is no gold standard for validation of the technique. Use of DCE-MRI to detect a reduction in perfusion related to hypoxia in the menstrual cycle would require a specialised imaging protocol and robust data analysis using a complex model, including estimation of parameter uncertainties.

Other existing MRI techniques could be applied to measure endometrial hypoxia (**Fig. 2**). T2* is a characteristic tissue relaxation time that depends on inhomogeneities in the main magnetic field produced by the scanner as well as rapidly-changing inhomogeneities induced by the presence of other nearby molecules. Detection of a reduction in T2* is commonly assumed to be due to the presence of deoxyhaemoglobin and therefore tissue hypoxia. This technique has been used in the myometrium (Kido *et al.*, 2007; Imaoka *et al.*, 2012) and has the high spatial resolution necessary to investigate the endometrium. T2* can change for a number of other reasons, (e.g. local haematocrit, hemosiderin, calcification and tissue iron deposition) therefore changes should be interpreted with caution. Similarly, a non-invasive perfusion technique known as arterial spin labelling (ASL) (Ferré *et al.*, 2013) could be extended from existing work in the myometrium (Takahashi *et al.*, 2016) to the endometrium, though it can be technically challenging. Finally, the extensive work on hypoxia measurements in cancer (Horsman *et al.*, 2012) could be applied in the endometrium. Oxygen-

enhanced (OE) MRI (O'Connor, Robinson & Waterton, 2019) allows a change in the tissue relaxation time T1 as a result of the patient breathing 100% oxygen through a mask to be related to the oxygen status of the tissue (O'Connor *et al.*, 2016). These minimally invasive MRI techniques may provide key information on the presence of human endometrial hypoxia throughout the menstrual cycle, with potential diagnostic and therapeutic benefits for women.

ROLE OF HYPOXIA THROUGHOUT THE MENSTRUAL CYCLE

Mice have the experimental advantage of genetic or pharmacological alteration to assess the role of hypoxia in endometrial function. HIF-1 α heterozygote mice have revealed that HIF-1 α is required for normal menstruation, and decreased HIF-1 α delays endometrial repair (Maybin *et al.*, 2018). Pharmacological stabilisation and inhibition of HIF-1 α in mice has confirmed this role (Chen *et al.*, 2015; Maybin *et al.*, 2018). Mice placed in hyperoxic chambers (75% O₂) during menses had reduced local endometrial hypoxia at menstruation and delayed endometrial repair (Maybin *et al.*, 2018). HIF-2 α deficiency restricted to uterine stromal cells in a mouse implantation model revealed a key role in decidualisation, endometrial receptivity, embryonic implantation and survival (Matsumoto *et al.*, 2018).

This emerging evidence for the presence and important role of hypoxia and HIF in endometrial function presents an exciting and developing research area (**Fig. 3**). The effects of hypoxia on the important menstrual processes of inflammation, proliferation and tissue remodelling remains to be elucidated.

Impact of hypoxia on inflammation

Inflammation is a key event during implantation, at menstruation and the subsequent endometrial repair. There is a peri-menstrual influx of leukocytes into the endometrium, in particular neutrophils and macrophages (Armstrong *et al.*, 2017). Interactions between the inflammatory response and hypoxia are well described at other tissue sites (Cramer *et al.*, 2003; Taylor, 2008; Taylor *et al.*, 2016) but the impact of hypoxia on the endometrial inflammatory response is less well characterised.

Impact on neutrophils

Neutrophils comprise up to 15% of the total endometrial cell numbers during menstruation (Poropatich, Rojas & Silverberg, 1987; Salamonsen & Lathbury, 2000). Their influx is tightly regulated, displaying a rapid, short lasting induction, which coincides with the upregulation of chemokines and cytokines. This temporal dynamic has been observed in both the mouse model of simulated menses and in human endometrial samples (Armstrong *et al.*, 2017). Neutrophils are important mediators of endometrial breakdown, which has been confirmed by their depletion in the mouse model of menstruation (Kaitu'u-Lino, Morison & Salamonsen, 2007b). However, the depleting agent used in this study also affects the monocytic cell lineage. Activated neutrophils release enzymes such as neutrophil elastase and cathepsin G. These enzymes activate MMPs produced by endometrial stromal cells and cause degradation of the extracellular matrix (Salamonsen & Lathbury, 2000). In airway inflammation, hypoxia boosts neutrophil degranulation and protease release (Hoenderdos *et al.*, 2016). It would be informative to determine whether hypoxia has similar effects in the endometrial environment during menses.

Neutrophils also produce reactive oxygen species (ROS) that might participate in endometrial breakdown. The potential role of ROS in menstruation has been reported (Sugino *et al.*, 1996), suggesting that free oxygen radicals may contribute to endometrial shedding by causing tissue

407 damage. Indeed, the inhibition of ROS generation in the mouse model of simulated menstruation
408 has been shown to abrogate endometrial breakdown (Wu *et al.*, 2014).

409
410 Neutrophil depletion in mouse models also affected endometrial regeneration (Kaitu'u-Lino,
411 Morison & Salamonsen, 2007b). Little is known about the impact of hypoxia on neutrophils during
412 endometrial repair. The concept that hypoxia has an effect on neutrophil number and function is
413 derived from studies of tumour biology. In a mouse model of endometrial carcinoma there was
414 spatiotemporal correlation between hypoxia and neutrophil infiltration within the tumour (Blaisdell
415 *et al.*, 2015). Accumulation of pimonidazole and nuclear staining of HIF-1 α was detected slightly
416 prior to neutrophil infiltration. These results are consistent with those observed in the mouse model
417 of simulated menses, where pharmacological inhibition of HIF-1 α decreased the number of
418 endometrial neutrophils present during active bleeding (Maybin *et al.*, 2018). The role of hypoxia in
419 promoting neutrophil recruitment in endometrial carcinoma was confirmed by placing mice in
420 hyperoxic chambers (60% O₂) (Mahiddine *et al.*, 2019). This resulted in a dramatic reduction in
421 neutrophil influx within the tumour and also improved the ability of these cells to oppose tumour
422 growth through increased activation and expression of several MMPs and ROS production. This is
423 consistent with hypoxia not only affecting the recruitment of neutrophils, but also their function.
424 Determining the effects of hypoxia on neutrophil number and phenotype in the normal
425 endometrium would be of great interest to advance our understanding of menstrual physiology.

426
427 Effects of hypoxia on neutrophils have also been observed in benign tissues. Airway inflammation
428 studies have revealed that hypoxia, via HIF-1 α and HIF-2 α , prolonged neutrophil lifespan by
429 inhibiting apoptosis (Walmsley *et al.*, 2005; Thompson *et al.*, 2014). Glucocorticoids have also been
430 shown to delay neutrophil apoptosis *in vitro*, but this did not occur in the presence of hypoxia
431 (Marwick *et al.*, 2013). Neutrophil apoptosis has been identified in the menstrual endometrium of
432 mice, when hypoxia is present (Armstrong *et al.*, 2017). In addition, glucocorticoids have been

identified as having an important role in the human menstrual endometrium (McDonald *et al.*, 2006; Rae *et al.*, 2009). The impact of hypoxia on endometrial myeloid apoptosis has not been examined to date.

Impact on macrophages

Macrophages have been detected in the endometrium throughout the menstrual cycle, both close to the endometrial glands and in the stromal compartment (Bonatz *et al.*, 1992). They show a perimenstrual peak in number, reaching up to 15% of the cell total number at the time of menses (Salamonsen & Woolley, 1999). Like neutrophils, it is proposed that macrophages play a critical role in the onset of endometrial breakdown via production and release of MMPs (reviewed in (Critchley *et al.*, 2001; Thiruchelvam *et al.*, 2013)). There are also indications of their involvement in glandular remodelling (Garry *et al.*, 2010) and endometrial regeneration (Maybin *et al.*, 2012; Cousins, Kirkwood, *et al.*, 2016), including the regulation of angiogenesis (Thiruchelvam *et al.*, 2016).

Macrophages are remarkably plastic cells, capable of shifting towards different phenotypes by sensing the surrounding microenvironment (Martinez, 2008). Thus, their microenvironment may affect their recruitment and function. Historically, macrophage polarisation has been categorised as classical (M1) or alternative (M2). M1 phenotype is associated with microbicidal properties and M2 reflects a more regulatory, anti-inflammatory phenotype. More recently, macrophage polarisation is understood to be a dynamic spectrum of macrophage transition in response to environmental cues (Martinez & Gordon, 2014). As there is mounting evidence for hypoxia in the local endometrial environment at menstruation (Cousins, Murray, *et al.*, 2016; Maybin *et al.*, 2018), it is important to determine its effect on endometrial macrophages.

Under physiological conditions M2 macrophages are involved in angiogenesis and cellular clearance, hence promote wound healing. However, tumour-infiltrating macrophages (TAMs) are often correlated with poor cancer prognosis (Kawanaka *et al.*, 2008). TAMs have been shown to be retained in hypoxic regions of tumours through the Sema3A/Neuropilin-1 signaling axis, which is regulated by HIF-2 α (Casazza *et al.*, 2013). The influence of hypoxia on TAMs is not only limited to macrophage number but also influences their phenotype. Indeed, specific TAM phenotypical subsets have been reported depending on intra-tumoral oxygen levels (Laoui *et al.*, 2014).

Non-tumoral studies have also linked HIF to changes in macrophage phenotype. In a model of endotoxemia, HIF-1 α and HIF-2 α were differentially expressed in M1 and M2-macrophages respectively (Takeda *et al.*, 2010). In addition, in the context of obesity and adipose tissue inflammation, HIF-1 α has been proven to promote inflammation and insulin resistance through M1 macrophage polarisation whereas HIF-2 α ameliorated the effects via M2-macrophage induction (reviewed in (Lin & Simon, 2016)). Interestingly, HIF-1 α was found to be decreased in mouse adipose tissue when glucocorticoid activation was suppressed, suggesting a crucial role of glucocorticoids in HIF-dependent macrophage polarisation (Chapman *et al.*, 2013). Thus, different research fields converge around the concept that HIF-1 α may be required for M1 polarization of macrophages, while HIF-2 α might promote M2 polarization.

The menstrual endometrium presents a unique model of transient, physiological hypoxia in which to study macrophage number and phenotype. HIF-2 α may have a role in the recruitment and function of macrophages during implantation, when endometrial HIF-2 α was found to be present (Maybin *et al.*, 2018). However, a recent study of mice with a targeted deletion of HIF-1 α in myeloid cells resulted in decreased pregnancy rates and increased miscarriage rates, suggesting that HIF-1 α dependent pathways in myeloid cells are also important for maintenance of pregnancy (Köstlin-Gille

et al., 2019). It would be informative to establish if the balance between HIF-1 α /HIF-2 α determines the pro-inflammatory or anti-inflammatory fate of the endometrium.

Impact of hypoxia on proliferation

After 'injury', fibroblasts must migrate and proliferate in the damaged area, where they produce extracellular matrix (ECM) components that contribute to repair (Gonzalez *et al.*, 2016). This production must be tightly regulated to prevent excessive ECM growth, scar formation and fibrosis (Ruthenborg *et al.*, 2014). In dermal tissue, hypoxia has been shown to stimulate macrophage growth factors that may contribute to fibroblast proliferation and tissue repair (Murdoch, Muthana & Lewis, 2005). Macrophage production of platelet-derived growth factor (PDGF) enhances fibroblast mitosis, while transforming growth factor β (TGF- β) promotes the formation of the ECM (Ruthenborg *et al.*, 2014). In addition, hypoxia has been proven to induce the transcription of VEGF, connective tissue growth factor and adrenomedullin in endometrial stromal tissue (Maybin, Battersby, *et al.*, 2011; Maybin *et al.*, 2012). Hence, hypoxia may induce a pro-repair environment by modifying the secretome of endometrial cell populations.

To complete tissue restoration, reepithelialisation of the affected area must take place. In the skin, this is achieved through the migration and proliferation of keratinocytes towards the injury site (Ruthenborg *et al.*, 2014). Stabilisation of HIF-1 α in a mouse model of skin wound healing revealed its role in promoting keratinocyte proliferation and migration to the injured area, accelerating wound closure (Kalucka *et al.*, 2013). This is consistent with the findings of delayed endometrial repair with decreased HIF-1 α (Maybin *et al.*, 2018).

Impact of hypoxia on vascular remodelling and angiogenesis

Angiogenesis and vascular remodelling are crucial events in the endometrium throughout the menstrual cycle. Optimal vascular function is necessary to support the repair of the functional endometrial layer and to supply the thickened endometrium required for successful implantation and placentation.

VEGF is a key mediator of both physiological and tumoral angiogenesis and may be induced by hypoxia (Carmeliet, 2005). VEGF mRNA and protein have been detected during all phases of the menstrual cycle, both in the stromal compartment and the glandular epithelium (Stephen Charnock-Jones *et al.*, 1993; Shifren *et al.*, 1996; Punyadeera, 2006) but was maximal during menses (Sharkey *et al.*, 2000; Graubert *et al.*, 2001; Maybin, Hirani, *et al.*, 2011). Studies in mouse and macaque models of menstruation have shown that blocking VEGF dramatically decreases reepithelialisation and new blood vessel formation in the endometrium (Fan *et al.*, 2008), consistent with an essential role for VEGF in endometrial angiogenesis and repair.

Hypoxia has been detected in the mouse model of simulated menses (Chen *et al.*, 2015; Cousins, Murray, *et al.*, 2016; Maybin *et al.*, 2018) and coincides with increased VEGF mRNA (Cousins, Murray, *et al.*, 2016). Hypoxia and VEGF have also been detected in human perimenstrual endometrial biopsies (Punyadeera, 2006) highlighting their possible interrelation. *In vitro* studies have also shown that subjecting endometrial and epithelial stromal cells to hypoxia increases VEGF mRNA and protein (Popovici *et al.*, 1999; Sharkey *et al.*, 2000; Graubert *et al.*, 2001) and that silencing of HIF-1 α abrogates this hypoxia-induced VEGF expression (Maybin, Hirani, *et al.*, 2011; Chen *et al.*, 2015). Through a chromatin immunoprecipitation (ChIP) assay, Chen *et al.* detected the direct binding of HIF-1 α to the VEGF promoter, which was maximal during endometrial breakdown of the mouse model of menses (Chen *et al.*, 2015). Inhibition of HIF-1 α using 2-methoxyestradiol (2-

ME) significantly suppressed VEGF levels during menses. Therefore, hypoxia, and more specifically HIF-1 α , seems to promote endometrial VEGF during menses.

In addition, VEGF expression is induced by different cytokines and chemokines (Li *et al.*, 1995; Stavri *et al.*, 1995; Zagzag *et al.*, 2006), some of which contain hypoxic response elements. Optimal blood vessel formation requires the trafficking of endothelial progenitors cells through the interaction of the chemokine CXCL12 with its receptor CXCR4 (Ruthenborg *et al.*, 2014). Both ligand and receptor have been found to be upregulated by HIF-1 α , contributing to angiogenesis and blood vessel repair partly through VEGF (Ceradini *et al.*, 2004; Zagzag *et al.*, 2006). CXCL12 and CXCR4 have been described in the human endometrium (Ruiz *et al.*, 2010) and endometrial CXCR4 was found to be decreased in patients with heavy menstrual bleeding (Maybin *et al.*, 2018). Hence, the interactions between hypoxia pathways and inflammatory processes may significantly influence endometrial vascular function.

During decidualisation there is *in vitro* evidence that endometrial stromal cells increase VEGF mRNA and protein (Popovici *et al.*, 1999; Matsui *et al.*, 2004) and that hypoxia induced further increases in VEGF (Popovici *et al.*, 1999). This VEGF production may be responsible for macrophage recruitment and polarisation towards a pro-angiogenic M2 phenotype (Wheeler *et al.*, 2018). Thus, the responsiveness of the decidualised stroma to hypoxia suggests a possible role in the preparation of the endometrial vasculature for implantation. Uterine HIF2- α deficiency has been shown to impair decidualisation in mice, revealing a downregulation of prolactin-related factors which can compromise the maintenance of the corpus luteum and therefore endometrial receptivity (Matsumoto *et al.*, 2018).

When studying implantation in mice, HIF factors were found to be differentially expressed at the time of peri-implantation: HIF-1 α was detected in the luminal epithelium, whereas HIF-2 α

expression was limited to the stromal compartment and neither correlated with VEGF expression (Daikoku *et al.*, 2003). Therefore, HIF effects on implantation seem to be more versatile than simply contributing to vessel formation, playing a substantial role in decidualisation, endometrial receptivity and embryo survival (Matsumoto *et al.*, 2018). After implantation, HIF-1 α was found in the luminal epithelium and the decidual layer. However, the strongest signal came from HIF-2 α , whose expression was localised to stromal cells surrounding the blastocyst. This post-implantation HIF-2 α expression was correlated with VEGF induction, switching to a proangiogenic stimulus once implantation had taken place (Daikoku *et al.*, 2003).

THE ROLE OF HYPOXIA IN ENDOMETRIAL PATHOLOGY

As outlined above, the literature regarding the influence of hypoxia on inflammation, proliferation and vascular function is increasing (**Fig. 3**). The influence of oxygen levels on implantation, placentation and disorders such as pre-eclampsia has been comprehensively reviewed within this series by Burton *et al.* (Burton, 2009). The impact of hypoxia on embryo function has been covered in detail by Dunwoodie *et al.* (Dunwoodie, 2009). Therefore, this section is focused on the role of endometrial hypoxia during menstruation and its potential in the identification of novel diagnostic and therapeutic strategies.

Abnormal uterine bleeding

Abnormal uterine bleeding (AUB) affects 20-30% of pre-menopausal women and over 800,000 women seek treatment in the UK each year (**National Heavy Menstrual Bleeding Audit, 2011**). Available medical treatments are often discontinued due to side effects or lack of efficacy. Research in this area was previously hindered by lack of a consistent classification system for the diagnosis of

causes of AUB. This was rectified by the development of the FIGO classification system of structural and non-structural causes (Munro, Critchley & Fraser, 2011, 2018) (Fig. 4).

Structural causes of AUB

Structural causes of AUB can be detected on examination or imaging of the uterus, e.g polyps, adenomyosis, leiomyoma (fibroids) and malignancy (Munro, Critchley & Fraser, 2011, 2018). These conditions have previously been under-diagnosed, with clinicians often treating the symptom of AUB without identifying the underlying cause. This has limited our knowledge on why these conditions develop and why they result in AUB.

Adenomyosis is the presence of ectopic endometrial glands and stroma within the myometrial layer of the uterus. It occurs in 7-27% of reproductive aged women and presents with painful, heavy menstrual bleeding (Naftalin *et al.*, 2012; Mavrelos *et al.*, 2017). The impact of the adenomyotic lesions on the eutopic endometrium and the mechanisms causing AUB are not well understood. AUB due to adenomyosis (AUB-A) is particularly challenging as it is often resistant to medical treatment and surgical options (ablation or hysterectomy) are unacceptable to those wishing to preserve their fertility.

There is some evidence that the hypoxic response is aberrant within adenomyotic lesions. A study of hysterectomy samples from 14 women with adenomyosis and 9 without revealed increased VEGF protein in the eutopic endometrium of women with adenomyosis and increased VEGF and HIF-1 α protein in ectopic versus eutopic endometrium (Goteri *et al.*, 2009). This suggests that a hypoxic environment in the adenomyotic lesions could contribute to increased vessel formation. In endometriosis, where ectopic endometrium implants outside of the uterus, HIF-1 α was also found to be increased in ectopic versus eutopic endometrium (Wu *et al.*, 2007; Young *et al.*, 2014).

Inhibition of HIF-1 in a mouse model of endometriosis suppressed growth of lesions (Becker *et al.*, 2008), identifying the hypoxia pathway as a potential therapeutic target. The peritoneum is a common site for implantation of ectopic endometrial deposits in endometriosis. Women with endometriosis have been shown to have increased HIF-1 α in non-affected peritoneum compared to peritoneum from women without disease (Young *et al.*, 2014), consistent with a role of the hypoxia pathway in the development of peritoneal disease. Studies examining the non-affected myometrium in women with adenomyosis are not yet available, but similar alterations in hypoxic response would highlight hypoxia pathways as a potential target for preventative and therapeutic interventions.

Leiomyomas (uterine fibroids) are common, benign tumours of the myometrium that form as a consequence of the proliferation of uterine smooth muscle cells and collagen matrix. They occur in approximately 70% of women (Stewart *et al.*, 2017) and are extremely heterogeneous in size, location and pathophysiology. Leiomyoma are symptomatic in approximately 50% of women (Day Baird *et al.*, 2003) and may cause symptoms of AUB, pressure, pelvic pain and be associated with subfertility.

Genome wide association studies have identified genetic subgroups that may predispose to leiomyoma formation (reviewed in (Stewart *et al.*, 2016)) but local mechanisms regulating their development remain an area of active research. Uterine leiomyomas contain broad avascular areas and HIF-1 α protein was found to be increased in leiomyoma nuclear protein extracts when compared to adjacent myometrium (Ishikawa *et al.*, 2019). However, it is not yet clear whether hypoxia is necessary for leiomyoma development and/or growth. In contrast, an *in vivo* study of women with leiomyoma using DCE-MRI has revealed increased K^{trans} (a combination of perfusion and permeability) in fibroids compared with normal uterus (Majd *et al.*, 2017) which does not support the presence of hypoxia within fibroids. There is evidence that treatment of leiomyomas with gonadotrophin releasing hormone (GnRH) analogues, often used pre-operatively to reduce fibroid

size and decrease AUB, lead to a decrease in perfusion parameters (Munro *et al.*, 2014). These contrasting *in vitro* and *in vivo* findings may reflect the heterogeneity of leiomyomas and it remains unclear if altered perfusion is associated with AUB.

The cause of AUB experienced by a proportion of women with leiomyomas is not understood. Vasoconstriction may be impaired at the time of menstruation in women with fibroids, with leiomyoma tissue expressing altered levels of endothelin receptors and prostaglandin F2 α when compared to normal myometrium (Pekonen, Nyman & Rutanen, 1994; Miura *et al.*, 2006). A small decrease in spiral arteriole vasoconstriction can significantly increase menstrual blood flow, causing heavy menstrual bleeding. A greater understanding of the role of hypoxia in leiomyoma formation and growth may identify new, specific treatments to reduce their presence, size and symptoms.

Endometrial cancer. The importance of hypoxia in the tumour microenvironment is well established, including its influence on immune cell populations, angiogenesis, tumour progression and metastasis (De Bock, Mazzone & Carmeliet, 2011; Casazza *et al.*, 2014; Schito & Semenza, 2016; Semenza, 2016). The accuracy of translation of these principles to patients with endometrial cancer is less well determined. In a quest to identify a robust biomarker that would predict tumour behaviour, Chang *et al* identified an eight gene set of lymphocyte and tumour hypoxia markers and validated its performance in predicting overall survival in six cancers, including 370 women with endometrial cancer (Chang, Forde & Lai, 2019). They found a superior performance over current tumour staging parameters, highlighting the importance of hypoxia in determining risk and aiding clinical decision making.

Assessment of endometrial tissues from 386 patients with endometrial carcinoma using CAIX as a hypoxia marker and CD34 to determine vascular density, revealed that patients with the presence of both hypoxia and high vascular density (16.4%) had reduced disease-specific survival and distant

disease-free survival (Reijnen *et al.*, 2019). *In vivo* imaging with DCE-MRI revealed that a poor prognosis was associated with low microvascular blood flow to the endometrial tumour (Haldorsen *et al.*, 2013, 2014; Berg *et al.*, 2016). This was thought to reflect disorganised angiogenesis with coexisting vascular proliferation and hypoxia. These studies highlight normalisation of the vasculature to limit hypoxia as a potential therapeutic target in endometrial cancer.

Non-structural causes of AUB

These non-structural disorders are not usually identified by routine pelvic imaging. They include coagulopathies, ovulatory dysfunction, endometrial and iatrogenic causes (Munro, Critchley & Fraser, 2011, 2018). Evidence for a role of hypoxia in these disorders is limited but its contribution to AUB of endometrial origin (AUB-E) is discussed below.

AUB-E includes disorders of local endometrial haemostasis, vascular function and/or inflammation (**Fig. 4**). Women with objectively defined HMB (>80ml/cycle) had reduced levels of HIF-1 α protein and downstream target genes in menstrual phase endometrial biopsies when compared to those from women with normal blood loss (Maybin *et al.*, 2018). Examination of endometrial repair in mice where hypoxia was prevented during simulated menses, or where HIF-1 α was pharmacologically or genetically reduced, revealed delayed repair (Maybin *et al.*, 2018). This is consistent with hypoxia having a key role in the rapid endometrial repair necessary to limit menstrual blood loss. The delayed repair in a non-hypoxic mouse menstruation model could be rescued with a pharmacological compound that stabilises HIF-1, identifying a potential non-hormonal therapeutic target for women with AUB-E.

The cause of the endometrial tissue hypoxia observed at menstruation is unknown. It is likely that spiral arteriole vasoconstriction limits blood supply to the functional layer of the endometrium

following progesterone withdrawal (Markee, 1940). Hence, factors that limit the ability of the specialised endometrial arterioles to constrict will have a significant impact on the presence of endometrial hypoxia. Women with the symptom of HMB have been shown to have significantly decreased smooth muscle myosin heavy chain in their spiral arterioles and also reduced vascular smooth muscle cell proliferation during the mid-late secretory phase compared to those with normal menstrual blood loss (Abberton *et al.*, 1999). Another study showed that endometrial vessel wall circumference and endothelial cell focal discontinuities were both significantly larger in women with HMB compared to normal controls (Mints *et al.*, 2007). Furthermore, calponin (a vascular smooth muscle cell contractile protein) was found to be significantly lower in endometrial blood vessels in women with HMB (Biswas Shivhare *et al.*, 2014). This evidence is all consistent with an aberrant vasculature within the pre-menstrual endometrium of women with AUB-E, leading to a suboptimal hypoxic response during menstruation.

CONCLUSIONS

Herein, we have reviewed the mounting evidence for the presence of endometrial hypoxia and its potential impact on endometrial function. Furthering our understanding of hypoxia in endometrial physiology and pathology using the tools described in this review may provide novel preventative and therapeutic strategies for those suffering from endometrial disorders, including abnormal uterine bleeding (AUB). Furthermore, a complete understanding of optimal endometrial physiology may inform the management of other disorders where aberrant hypoxia is a prominent feature, such as tumour biology and chronic obstructive pulmonary disorder. Addressing the gaps in our knowledge of how hypoxia influences endometrial function represents an exciting area with huge translational potential.

ACKNOWLEDGEMENTS

We are grateful to Alison Murray for manuscript review and Sheila Milne for assistance with manuscript formatting. Some aspects of studies described herein were undertaken in the MRC Centre for Reproductive Health and have been supported by funding from the Medical Research Council: MRC Project Grant G0600048 and Centre Grants G1002033 and MR/N022556/1. This work was also supported by Wellcome Trust Grants 209589/Z/17/Z and 100646/Z/21/Z; Wellbeing of Women Grant RG1820; and Academy of Medical Sciences Grant AMS-SGCL13

Disclosure: RM-A, LK, JJR and JAM have nothing to disclose. HODC has clinical research support for laboratory consumables and staff from Bayer AG and provides consultancy advice (but with no personal remuneration) for Bayer AG, PregLem SA, Gedeon Richter, Vifor Pharma UK Ltd, AbbVie Inc; Myovant Sciences GmbH. HODC receives royalties from UpToDate for article on abnormal uterine bleeding. Jackie Maybin is a Guest Editor of this special issue of Reproduction. Jackie Maybin was not involved in the review or editorial process for this paper, on which she is listed as an author.

Author contributions: RM-A, LK and JAM wrote and edited the manuscript. JJR and HODC modified the manuscript. All authors have read and approved the final version of this manuscript.

728 **REFERENCES**

729 Abberton KM, Taylor NH, Healy DL & Rogers PAW 1999 Vascular smooth muscle cell proliferation in
730 arterioles of the human endometrium. *Human Reproduction* **14** 1072–1079.

731 Armstrong GM, Maybin JA, Murray AA, Nicol M, Walker C, Saunders PTK, Rossi AG & Critchley HOD
732 2017 Endometrial apoptosis and neutrophil infiltration during menstruation exhibits spatial and
733 temporal dynamics that are recapitulated in a mouse model. *Scientific Reports* **7** 1–14.

734 Baird DT, Cameron ST, Critchley HOD, Drudy TA, Howe A, Jones RL, Lea RG & Kelly RW 1996
735 Prostaglandins and menstruation. *European Journal of Obstetrics & Gynecology and Reproductive*
736 *Biology* **70** 15–17.

737 Becker CM, Rohwer N, Funakoshi T, Cramer T, Bernhardt W, Birsner A, Folkman J & D’Amato RJ 2008
738 2-methoxyestradiol inhibits hypoxia-inducible factor-1 α and suppresses growth of lesions in a mouse
739 model of endometriosis. *The American Journal of Pathology* **172** 534–544.

740 Bellofiore N, Ellery SJ, Mamrot J, Walker DW, Temple-Smith P & Dickinson H 2017 First evidence of a
741 menstruating rodent: the spiny mouse (*Acomys cahirinus*). *American Journal of Obstetrics and*
742 *Gynecology* **216** 40.e1-40.e11.

743 Bellofiore N, Rana S, Dickinson H, Temple-Smith P & Evans J 2018 Characterization of human-like
744 menstruation in the spiny mouse: comparative studies with the human and induced mouse model.
745 *Human Reproduction* **33** 1715–1726.

746 Berg A *et al.* 2016 Tissue and imaging biomarkers for hypoxia predict poor outcome in endometrial
747 cancer. *Oncotarget* **7** 69844–69856.

748 Biswas Shivhare S, Bulmer JN, Innes BA, Hapangama DK & Lash GE 2014 Altered vascular smooth
749 muscle cell differentiation in the endometrial vasculature in menorrhagia. *Human Reproduction* **29**
750 1884–1894.

751 Blaisdell A, Crequer A, Columbus D, Daikoku T, Mittal K, Dey SK & Erlebacher A 2015 Neutrophils
752 oppose uterine epithelial carcinogenesis via debridement of hypoxic tumor cells. *Cancer Cell* **28**
753 785–799.

754 De Bock K, Mazzone M & Carmeliet P 2011 Antiangiogenic therapy, hypoxia, and metastasis: risky
755 liaisons, or not? *Nature Reviews Clinical Oncology* **8** 393–404.

756 Bonatz G, Hansmann M-L, Buchholz F, Mettler L, Radzun HJ & Semm K 1992 Macrophage- and

- 757 lymphocyte-subtypes in the endometrium during different phases of the ovarian cycle. *International*
 758 *Journal of Gynecology & Obstetrics* **37** 29–36.
- 759 Brasted M, White CA, Kennedy TG & Salamonsen LA 2003 Mimicking the events of menstruation in
 760 the murine uterus. *Biology of Reproduction* **69** 1273–1280.
- 761 Brenner RM & Slayden OD 2012 Molecular and functional aspects of menstruation in the macaque.
 762 *Reviews in Endocrine and Metabolic Disorders* **13** 309–318.
- 763 Brix G, Kiessling F, Lucht R, Darai S, Wasser K, Delorme S & Griebel J 2004 Microcirculation and
 764 microvasculature in breast tumors: pharmacokinetic analysis of dynamic MR image series. *Magnetic*
 765 *Resonance in Medicine* **52** 420–429.
- 766 Brix G, Zwick S, Kiessling F & Griebel J 2009 Pharmacokinetic analysis of tissue microcirculation using
 767 nested models: multimodel inference and parameter identifiability. *Medical Physics* **36** 2923–2933.
- 768 Burton GJ 2009 Oxygen, the Janus gas; its effects on human placental development and function.
 769 *Journal of Anatomy* **215** 27–35.
- 770 Carmeliet P 2005 VEGF as a key mediator of angiogenesis in cancer. *Oncology* **69** 4–10.
- 771 Casazza A *et al.* 2013 Impeding macrophage entry into hypoxic tumor areas by Sema3A/Nrp1
 772 signaling blockade inhibits angiogenesis and restores antitumor immunity. *Cancer Cell* **24** 695–709.
- 773 Casazza A, Di Conza G, Wenes M, Finisguerra V, Deschoemaeker S & Mazzone M 2014 Tumor
 774 stroma: a complexity dictated by the hypoxic tumor microenvironment. *Oncogene* **33** 1743–1754.
- 775 Celik O, Hascalik S, Sarac K, Meydanli MM, Alkan A & Mizrak B 2005 Magnetic resonance
 776 spectroscopy of premalignant and malignant endometrial disorders: a feasibility of in vivo study.
 777 *European Journal of Obstetrics & Gynecology and Reproductive Biology* **118** 241–245.
- 778 Ceradini DJ *et al.* 2004 Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1
 779 induction of SDF-1. *Nature Medicine* **10** 858–864.
- 780 Chang WH, Forde D & Lai AG 2019 A novel signature derived from immunoregulatory and hypoxia
 781 genes predicts prognosis in liver and five other cancers. *Journal of Translational Medicine* **17** 14.
- 782 Chapman KE, Coutinho AE, Zhang Z, Kipari T, Savill JS & Seckl JR 2013 Changing glucocorticoid action:
 783 11 β -Hydroxysteroid dehydrogenase type 1 in acute and chronic inflammation. *The Journal of Steroid*
 784 *Biochemistry and Molecular Biology* **137** 82–92.
- 785 Chen X *et al.* 2015 Vascular endothelial growth factor (VEGF) regulation by hypoxia inducible factor-1

- 786 alpha (HIF1A) starts and peaks during endometrial breakdown, not repair, in a mouse menstrual-like
787 model. *Human Reproduction* **30** 2160–2170.
- 788 Chen X *et al.* 2020 Hypoxia: involved but not essential for endometrial breakdown in mouse
789 menstrual-like model. *Reproduction* **159** 133–144.
- 790 Corker CS, Michie E, Hobson B & Parboosingh J 1976 Hormonal patterns in conceptual cycles and
791 early pregnancy. *British Journal of Obstetrics and Gynaecology* **83** 489–494.
- 792 Coudyzer P, Lemoine P, Jordan BF, Gallez B, Galant C, Nisolle M, Courtoy PJ, Henriët P & Marbaix E
793 2013 Hypoxia is not required for human endometrial breakdown or repair in a xenograft model of
794 menstruation. *The FASEB Journal* **27** 3711–3719.
- 795 Coudyzer P, Lemoine P, Po C, Jordan BF, Van Der Smissen P, Courtoy PJ, Henriët P & Marbaix E 2015
796 Induction of post-menstrual regeneration by ovarian steroid withdrawal in the functionalis of
797 xenografted human endometrium. *Human Reproduction* **30** 1156–1168.
- 798 Cousins FL, Kirkwood PM, Saunders PTK & Gibson DA 2016 Evidence for a dynamic role for
799 mononuclear phagocytes during endometrial repair and remodelling. *Scientific Reports* **6** 36748.
- 800 Cousins FL, Murray A, Esnal A, Gibson DA, Critchley HOD & Saunders PTK 2014 Evidence from a
801 mouse model that epithelial cell migration and mesenchymal-epithelial transition contribute to rapid
802 restoration of uterine tissue integrity during menstruation. *PLoS ONE* **9** e86378.
- 803 Cousins FL, Murray AA, Scanlon JP & Saunders PTK 2016 Hypoxyprobe™ reveals dynamic spatial and
804 temporal changes in hypoxia in a mouse model of endometrial breakdown and repair. *BMC Research*
805 *Notes* **9** 30.
- 806 Cramer T *et al.* 2003 HIF-1 α is essential for myeloid cell-mediated inflammation. *Cell* **112** 645–657.
- 807 Critchley HOD, Henderson TA, Kelly RW, Scobie GS, Evans LR, Groome NP & Saunders PTK 2002 Wild-
808 type estrogen receptor (ER β 1) and the splice variant (ER β cx/ β 2) are both expressed within the
809 human endometrium throughout the normal menstrual cycle. *The Journal of Clinical Endocrinology*
810 *& Metabolism* **87** 5265–5273.
- 811 Critchley HOD, Jones RL, Lea RG, Drudy TA, Kelly RW, Williams ARW & Baird DT 1999 Role of
812 inflammatory mediators in human endometrium during progesterone withdrawal and early
813 pregnancy. *The Journal of Clinical Endocrinology & Metabolism* **84** 240–248.
- 814 Critchley HOD, Kelly RW, Brenner RM & Baird DT 2001 The endocrinology of menstruation - a role
815 for the immune system. *Clinical Endocrinology* **55** 701–710.

- 816 Critchley HOD, Maybin JA, Armstrong GM & Williams ARW 2020 Physiology of the endometrium and
817 regulation of menstruation. *Physiological Reviews* **100** 1149–1179.
- 818 Critchley HOD, Osei J, Henderson TA, Boswell L, Sales KJ, Jabbour HN & Hirani N 2006 Hypoxia-
819 inducible factor-1 α expression in human endometrium and its regulation by prostaglandin E-series
820 prostanoid receptor 2 (EP2). *Endocrinology* **147** 744–753.
- 821 Daikoku T, Matsumoto H, Gupta RA, Das SK, Gassmann M, DuBois RN & Dey SK 2003 Expression of
822 hypoxia-inducible factors in the peri-implantation mouse uterus is regulated in a cell-specific and
823 ovarian steroid hormone-dependent manner. *Journal of Biological Chemistry* **278** 7683–7691.
- 824 Day Baird D, Dunson DB, Hill MC, Cousins D & Schectman JM 2003 High cumulative incidence of
825 uterine leiomyoma in black and white women: ultrasound evidence. *American Journal of Obstetrics
826 and Gynecology* **188** 100–107.
- 827 Donoghue JF, McGavigan CJ, Lederman FL, Cann LM, Fu L, Dimitriadis E, Girling JE & Rogers PAW
828 2012 Dilated thin-walled blood and lymphatic vessels in human endometrium: a potential role for
829 VEGF-D in progestin-induced break-through bleeding. *PLoS ONE* **7** e30916.
- 830 Dunn CL, Kelly RW & Critchley HO 2003 Decidualization of the human endometrial stromal cell: an
831 enigmatic transformation. *Reproductive BioMedicine Online* **7** 151–161.
- 832 Dunwoodie SL 2009 The role of hypoxia in development of the mammalian embryo. *Developmental
833 Cell* **17** 755–773.
- 834 Evans J, Kaitu'u-Lino T & Salamonsen LA 2011 Extracellular matrix dynamics in scar-free endometrial
835 repair: perspectives from mouse in vivo and human in vitro studies. *Biology of Reproduction* **85** 511–
836 523.
- 837 Fan X, Krieg S, Kuo CJ, Wiegand SJ, Rabinovitch M, Druzin ML, Brenner RM, Giudice LC & Nayak NR
838 2008 VEGF blockade inhibits angiogenesis and reepithelialization of endometrium. *The FASEB
839 Journal* **22** 3571–3580.
- 840 Ferré J-C, Bannier E, Raoult H, Mineur G, Carsin-Nicol B & Gauvrit J-Y 2013 Arterial spin labeling (ASL)
841 perfusion: techniques and clinical use. *Diagnostic and Interventional Imaging* **94** 1211–1223.
- 842 Finn CA 1986 Implantation, menstruation and inflammation. *Biological Reviews* **61** 313–328.
- 843 Finn CA & Pope M 1984 Vascular and cellular changes in the decidualized endometrium of the
844 ovariectomized mouse following cessation of hormone treatment: a possible model for
845 menstruation. *Journal of Endocrinology* **100** 295–300.

- 846 Fraser IS, McCarron G, Hutton B & Macey D 1987 Endometrial blood flow measured by xenon 133
 847 clearance in women with normal menstrual cycles and dysfunctional uterine bleeding. *American*
 848 *Journal of Obstetrics and Gynecology* **156** 158–166.
- 849 Gaide Chevrionnay HP, Galant C, Lemoine P, Courtoy PJ, Marbaix E & Henriët P 2009 Spatiotemporal
 850 coupling of focal extracellular matrix degradation and reconstruction in the menstrual human
 851 endometrium. *Endocrinology* **150** 5094–5105.
- 852 Gannon BJ, Carati CJ & Verco CJ 1997 Endometrial perfusion across the normal human menstrual
 853 cycle assessed by laser Doppler fluxmetry. *Human Reproduction* **12** 132–139.
- 854 Garry R, Hart R, Karthigasu K & Burke C 2010 Structural changes in endometrial basal glands during
 855 menstruation. *BJOG: An International Journal of Obstetrics & Gynaecology* **117** 1175–1185.
- 856 Gellersen B & Brosens JJ 2014 Cyclic decidualization of the human endometrium in reproductive
 857 health and failure. *Endocrine Reviews* **35** 851–905.
- 858 Gonzalez AC de O, Costa TF, Andrade Z de A & Medrado ARAP 2016 Wound healing - A literature
 859 review. *Anais brasileiros de dermatologia* **91** 614–620.
- 860 Goteri G, Lucarini G, Montik N, Zizzi A, Stramazotti D, Fabris G, Tranquilli AL & Ciavattini A 2009
 861 Expression of vascular endothelial growth factor (VEGF), hypoxia inducible factor-1 α (HIF-1 α), and
 862 microvessel density in endometrial tissue in women with adenomyosis. *International Journal of*
 863 *Gynecological Pathology* **28** 157–163.
- 864 Graubert MD, Asuncion Ortega M, Kessel B, Mortola JF & Iruela-Arispe ML 2001 Vascular repair after
 865 menstruation involves regulation of vascular endothelial growth factor-receptor phosphorylation by
 866 sFLT-1. *The American Journal of Pathology* **158** 1399–1410.
- 867 Guo Y, He B, Xu X & Wang J 2011 Comprehensive analysis of leukocytes, vascularization and matrix
 868 metalloproteinases in human menstrual xenograft model. *PLoS ONE* **6** e16840.
- 869 Haldorsen IS *et al.* 2013 Dynamic contrast-enhanced MRI in endometrial carcinoma identifies
 870 patients at increased risk of recurrence. *European Radiology* **23** 2916–2925.
- 871 Haldorsen IS *et al.* 2014 Increased microvascular proliferation is negatively correlated to tumour
 872 blood flow and is associated with unfavourable outcome in endometrial carcinomas. *British Journal*
 873 *of Cancer* **110** 107–114.
- 874 Hamlett GWD 1934 Uterine bleeding in a bat, *Glossophaga soricina*. *The Anatomical Record* **60** 9–
 875 17.

- 876 Hoenderdos K *et al.* 2016 Hypoxia upregulates neutrophil degranulation and potential for tissue
877 injury. *Thorax* **71** 1030–1038.
- 878 Horsman MR, Mortensen LS, Petersen JB, Busk M & Overgaard J 2012 Imaging hypoxia to improve
879 radiotherapy outcome. *Nature Reviews Clinical Oncology* **9** 674–687.
- 880 Van der Horst CJ & Gillman J 1940 Mechanism of ovulation and corpus luteum formation in
881 *Elephantulus*. *Nature* **145** 974–974.
- 882 Imaoka I *et al.* 2012 T2* relaxometry mapping of the uterine zones. *Acta Radiologica* **53** 473–477.
- 883 Ishikawa H, Xu L, Sone K, Kobayashi T, Wang G & Shozu M 2019 Hypoxia induces hypoxia-inducible
884 factor 1 α and potential HIF-responsive gene expression in uterine leiomyoma. *Reproductive Sciences*
885 **26** 428–435.
- 886 Kaitu'u-Lino TJ, Morison NB & Salamonsen LA 2007a Estrogen is not essential for full endometrial
887 restoration after breakdown: lessons from a mouse model. *Endocrinology* **148** 5105–5111.
- 888 Kaitu'u-Lino TJ, Morison NB & Salamonsen LA 2007b Neutrophil depletion retards endometrial repair
889 in a mouse model. *Cell and Tissue Research* **328** 197–206.
- 890 Kalucka J *et al.* 2013 Loss of epithelial hypoxia-inducible factor prolyl hydroxylase 2 accelerates skin
891 wound healing in mice. *Molecular and Cellular Biology* **33** 3426–3438.
- 892 Kawanaka T, Kubo A, Ikushima H, Sano T, Takegawa Y & Nishitani H 2008 Prognostic significance of
893 HIF-2 α expression on tumor infiltrating macrophages in patients with uterine cervical cancer
894 undergoing radiotherapy. *The Journal of Medical Investigation* **55** 78–86.
- 895 Kelly R, King A & Critchley H 2001 Cytokine control in human endometrium. *Reproduction* **121** 3–19.
- 896 Kido A, Koyama T, Kataoka M, Yamamoto A, Saga T, Turner R & Togashi K 2007 Physiological changes
897 of the human uterine myometrium during menstrual cycle: preliminary evaluation using BOLD MR
898 imaging. *Journal of Magnetic Resonance Imaging* **26** 695–700.
- 899 King AE, Critchley HO & Kelly RW 2001 The NF-kappaB pathway in human endometrium and first
900 trimester decidua. *Molecular Human Reproduction* **7** 175–183.
- 901 Köstlin-Gille N, Dietz S, Schwarz J, Spring B, Pauluschke-Fröhlich J, Poets CF & Gille C 2019 HIF-1 α -
902 deficiency in myeloid cells leads to a disturbed accumulation of myeloid derived suppressor cells
903 (MDSC) during pregnancy and to an increased abortion rate in mice. *Frontiers in Immunology* **10**
904 161.

- 905 Kuokkanen S, Zhu L & Pollard JW 2017 Xenografted tissue models for the study of human
906 endometrial biology. *Differentiation* **98** 62–69.
- 907 Kupesic S & Kurjak A 1993 Uterine and ovarian perfusion during the periovulatory period assessed by
908 transvaginal color Doppler. *Fertility and Sterility* **60** 439–443.
- 909 Lange T, Dydak U, Roberts TPL, Rowley HA, Bjeljac M & Boesiger P 2006 Pitfalls in lactate
910 measurements at 3T. *AJNR. American Journal of Neuroradiology* **27** 895–901.
- 911 Laoui D *et al.* 2014 Tumor hypoxia does not drive differentiation of tumor-associated macrophages
912 but rather fine-tunes the M2-like macrophage population. *Cancer Research* **74** 24–30.
- 913 Lessey BA, Killam AP, Metzger DA, Haney AF, Greene GL & McCarty KS 1988 Immunohistochemical
914 analysis of human uterine estrogen and progesterone receptors throughout the menstrual cycle. *The*
915 *Journal of Clinical Endocrinology & Metabolism* **67** 334–340.
- 916 Li J *et al.* 1995 Induction of vascular endothelial growth factor gene expression by interleukin-1 in rat
917 aortic smooth muscle cells. *Journal of Biological Chemistry* **270** 308–312.
- 918 Lin N & Simon MC 2016 Hypoxia-inducible factors: key regulators of myeloid cells during
919 inflammation. *Journal of Clinical Investigation* **126** 3661–3671.
- 920 Mahiddine K, Blaisdell A, Ma S, Créquer-Grandhomme A, Lowell CA & Erlebacher A 2019 Relief of
921 tumor hypoxia unleashes the tumoricidal potential of neutrophils. *Journal of Clinical Investigation*
922 **130** 389–403.
- 923 Majd S, Johan Jeet Abdullah B, Ahmad Sarji S, Hong Yeong C & Azmi Mat Adenan N 2017
924 Quantitative dynamic contrast-enhanced magnetic resonance imaging in symptomatic uterine
925 fibroids and normal uterus: a feasibility study. *Iranian Journal of Radiology* **14** 1–10.
- 926 Markee JE 1940 Menstruation in intraocular endometrial transplants in the Rhesus monkey. *Contrib.*
927 *Embryol.* 219–308.
- 928 Marsh MM, Malakooti N, Taylor NH, Findlay JK & Salamonsen LA 1997 Endothelin and neutral
929 endopeptidase in the endometrium of women with menorrhagia. *Human Reproduction* **12** 2036–
930 2040.
- 931 Martinez FO 2008 Macrophage activation and polarization. *Frontiers in Bioscience* **13** 453–461.
- 932 Martinez FO & Gordon S 2014 The M1 and M2 paradigm of macrophage activation: time for
933 reassessment. *F1000Prime Reports* **6**

- 934 Marwick JA *et al.* 2013 Oxygen levels determine the ability of glucocorticoids to influence neutrophil
935 survival in inflammatory environments. *Journal of Leukocyte Biology* **94** 1285–1292.
- 936 Masuda H *et al.* 2007 Noninvasive and real-time assessment of reconstructed functional human
937 endometrium in NOD/SCID/gamma c(null) immunodeficient mice. *Proceedings of the National*
938 *Academy of Sciences* **104** 1925–1930.
- 939 Matsui N, Kawano Y, Nakamura S & Miyakawa I 2004 Changes in vascular endothelial growth factor
940 production associated with decidualization by human endometrial stromal cells in vitro. *Acta*
941 *Obstetrica et Gynecologica Scandinavica* **83** 138–143.
- 942 Matsumoto L *et al.* 2018 HIF2 α in the uterine stroma permits embryo invasion and luminal
943 epithelium detachment. *Journal of Clinical Investigation* **128** 3186–3197.
- 944 Matsuura-Sawada R *et al.* 2005 Reproduction of menstrual changes in transplanted human
945 endometrial tissue in immunodeficient mice. *Human Reproduction* **20** 1477–1484.
- 946 Mavrellos D, Holland TK, O'Donovan O, Khalil M, Ploumpidis G, Jurkovic D & Khalaf Y 2017 The impact
947 of adenomyosis on the outcome of IVF–embryo transfer. *Reproductive BioMedicine Online* **35** 549–
948 554.
- 949 Maybin JA, Barcroft J, Thiruchelvam U, Hirani N, Jabbour HN & Critchley HOD 2012 The presence and
950 regulation of connective tissue growth factor in the human endometrium. *Human Reproduction* **27**
951 1112–1121.
- 952 Maybin JA, Battersby S, Hirani N, Nikitenko LL, Critchley HOD & Jabbour HN 2011 The expression and
953 regulation of adrenomedullin in the human endometrium: a candidate for endometrial repair.
954 *Endocrinology* **152** 2845–2856.
- 955 Maybin JA, Hirani N, Brown P, Jabbour HN & Critchley HOD 2011 The regulation of vascular
956 endothelial growth factor by hypoxia and prostaglandin F2 α during human endometrial repair. *The*
957 *Journal of Clinical Endocrinology & Metabolism* **96** 2475–2483.
- 958 Maybin JA, Murray AA, Saunders PTK, Hirani N, Carmeliet P & Critchley HOD 2018 Hypoxia and
959 hypoxia inducible factor-1 α are required for normal endometrial repair during menstruation. *Nature*
960 *Communications* **9** 295.
- 961 McDonald SE, Henderson TA, Gomez-Sanchez CE, Critchley HOD & Mason JI 2006 11 β -
962 Hydroxysteroid dehydrogenases in human endometrium. *Molecular and Cellular Endocrinology* **248**
963 72–78.

- 964 Michaely HJ, Sourbron SP, Buettner C, Lodemann K-P, Reiser MF & Schoenberg SO 2008 Temporal
 965 constraints in renal perfusion imaging with a 2-compartment model. *Investigative Radiology* **43**
 966 120–128.
- 967 Mints M, Hultenby K, Zetterberg E, Blomgren B, Falconer C, Rogers R & Palmblad J 2007 Wall
 968 discontinuities and increased expression of vascular endothelial growth factor-A and vascular
 969 endothelial growth factor receptors 1 and 2 in endometrial blood vessels of women with
 970 menorrhagia. *Fertility and Sterility* **88** 691–697.
- 971 Miura S, Khan KN, Kitajima M, Hiraki K, Moriyama S, Masuzaki H, Samejima T, Fujishita A & Ishimaru
 972 T 2006 Differential infiltration of macrophages and prostaglandin production by different uterine
 973 leiomyomas. *Human Reproduction* **21** 2545–2554.
- 974 Mole DR, Blancher C, Copley RR, Pollard PJ, Gleadle JM, Ragoussis J & Ratcliffe PJ 2009 Genome-wide
 975 association of hypoxia-inducible factor (HIF)-1 α and HIF-2 α DNA binding with expression profiling of
 976 hypoxia-inducible transcripts. *Journal of Biological Chemistry* **284** 16767–16775.
- 977 Munro KI *et al.* 2014 Quantitative serial MRI of the treated fibroid uterus. *PLoS ONE*. Edited by S.
 978 Franks **9** e89809.
- 979 Munro MG, Critchley HOD & Fraser IS 2011 The FIGO classification of causes of abnormal uterine
 980 bleeding in the reproductive years. *Fertility and Sterility* **95** 2204–2208.
- 981 Munro MG, Critchley HOD & Fraser IS 2018 The two FIGO systems for normal and abnormal uterine
 982 bleeding symptoms and classification of causes of abnormal uterine bleeding in the reproductive
 983 years: 2018 revisions. *International Journal of Gynecology & Obstetrics* **143** 393–408.
- 984 Murdoch C, Muthana M & Lewis CE 2005 Hypoxia regulates macrophage functions in inflammation.
 985 *The Journal of Immunology* **175** 6257–6263.
- 986 Mutsaers SE, Birnie K, Lansley S, Herrick SE, Lim C-B & Prêle CM 2015 Mesothelial cells in tissue
 987 repair and fibrosis. *Frontiers in Pharmacology* **6**
- 988 Naftalin J, Hoo W, Pateman K, Mavrelos D, Holland T & Jurkovic D 2012 How common is
 989 adenomyosis? A prospective study of prevalence using transvaginal ultrasound in a gynaecology
 990 clinic. *Human Reproduction* **27** 3432–3439.
- 991 Noyes RW, Hertig AT & Rock J 1950 Dating the endometrial biopsy. *Fertility and Sterility* **1** 3–25.
- 992 O'Connor JPB *et al.* 2016 Oxygen-enhanced MRI accurately identifies, quantifies, and maps tumor
 993 hypoxia in preclinical cancer models. *Cancer Research* **76** 787–795.

- 994 O'Connor JPB, Robinson SP & Waterton JC 2019 Imaging tumour hypoxia with oxygen-enhanced MRI
995 and BOLD MRI. *The British Journal of Radiology* **92** 20180642.
- 996 Pasanen A, Heikkilä M, Rautavuoma K, Hirsilä M, Kivirikko KI & Myllyharju J 2010 Hypoxia-inducible
997 factor (HIF)-3 α is subject to extensive alternative splicing in human tissues and cancer cells and is
998 regulated by HIF-1 but not HIF-2. *The International Journal of Biochemistry & Cell Biology* **42** 1189–
999 1200.
- 1000 Pekonen F, Nyman T & Rutanen E-M 1994 Differential expression of mRNAs for endothelin-related
1001 proteins in human endometrium, myometrium and leiomyoma. *Molecular and Cellular*
1002 *Endocrinology* **103** 165–170.
- 1003 Polotsky AJ, Zhu L, Santoro N & Pollard JW 2009 Lithium chloride treatment induces epithelial cell
1004 proliferation in xenografted human endometrium. *Human Reproduction* **24** 1960–1967.
- 1005 Popovici RM, Irwin JC, Giaccia AJ & Giudice LC 1999 Hypoxia and cAMP stimulate vascular
1006 endothelial growth factor (VEGF) in human endometrial stromal cells: potential relevance to
1007 menstruation and endometrial regeneration. *The Journal of Clinical Endocrinology & Metabolism* **84**
1008 2245–2245.
- 1009 Poropatich C, Rojas M & Silverberg SG 1987 Polymorphonuclear leukocytes in the endometrium
1010 during the normal menstrual cycle. *International Journal of Gynecological Pathology* **6** 230–234.
- 1011 Prill HJ & Götz F 1961 Blood flow in the myometrium and endometrium of the uterus. *American*
1012 *Journal of Obstetrics and Gynecology* **82** 102–108.
- 1013 Pringle KG, Kind KL, Thompson JG & Roberts CT 2007 Complex interactions between hypoxia
1014 inducible factors, insulin-like growth factor-II and oxygen in early murine trophoblasts. *Placenta* **28**
1015 1147–1157.
- 1016 Punyadeera C 2006 Expression and regulation of vascular endothelial growth factor ligands and
1017 receptors during menstruation and post-menstrual repair of human endometrium. *Molecular*
1018 *Human Reproduction* **12** 367–375.
- 1019 Rae M, Mohamad A, Price D, Hadoke PWF, Walker BR, Mason JI, Hillier SG & Critchley HOD 2009
1020 Cortisol inactivation by 11 β -hydroxysteroid dehydrogenase-2 may enhance endometrial
1021 angiogenesis via reduced thrombospondin-1 in heavy menstruation. *The Journal of Clinical*
1022 *Endocrinology & Metabolism* **94** 1443–1450.
- 1023 Raine-Fenning NJ 2004 Quantifying the changes in endometrial vascularity throughout the normal
1024 menstrual cycle with three-dimensional power Doppler angiography. *Human Reproduction* **19** 330–

- 1025 338.
- 1026 Rasweiler JJ & de Bonilla H 1992 Menstruation in short-tailed fruit bats (*Carollia* spp.). *Reproduction*
 1027 **95** 231–248.
- 1028 Reijnen C *et al.* 2019 Poor outcome in hypoxic endometrial carcinoma is related to vascular density.
 1029 *British Journal of Cancer* **120** 1037–1044.
- 1030 Royal College of Obstetricians and Gynaecologists 2014 *National Heavy Menstrual Bleeding Audit*
- 1031 Rudolph M, Döcke W-D, Müller A, Menning A, Röse L, Zollner TM & Gashaw I 2012 Induction of overt
 1032 menstruation in intact mice. *PLoS ONE* **7** e32922.
- 1033 Ruiz A, Salvo VA, Ruiz LA, Báez P, García M & Flores I 2010 Basal and steroid hormone-regulated
 1034 expression of CXCR4 in human endometrium and endometriosis. *Reproductive Sciences* **17** 894–903.
- 1035 Ruthenborg RJ, Ban J-J, Wazir A, Takeda N & Kim J 2014 Regulation of wound healing and fibrosis by
 1036 hypoxia and hypoxia-inducible factor-1. *Molecules and Cells* **37** 637–643.
- 1037 Salamonsen LA & Lathbury LJ 2000 Endometrial leukocytes and menstruation. *Human Reproduction*
 1038 *Update* **6** 16–27.
- 1039 Salamonsen LA & Woolley DE 1999 Menstruation: induction by matrix metalloproteinases and
 1040 inflammatory cells. *Journal of Reproductive Immunology* **44** 1–27.
- 1041 Salceda S & Caro J 1997 Hypoxia-inducible factor 1 α (HIF-1 α) protein is rapidly degraded by the
 1042 ubiquitin-proteasome system under normoxic conditions. *Journal of Biological Chemistry* **272**
 1043 22642–22647.
- 1044 Sarac K, Celik O, Hascalik S, Alkan A & Mizrak B 2004 In vivo proton magnetic resonance
 1045 spectroscopy in the evaluation of the endometrium. *Acta Obstetrica et Gynecologica Scandinavica*
 1046 **83** 751–757.
- 1047 Schito L & Semenza GL 2016 Hypoxia-inducible factors: master regulators of cancer progression.
 1048 *Trends in Cancer* **2** 758–770.
- 1049 Semenza GL 2000 HIF-1: mediator of physiological and pathophysiological responses to hypoxia.
 1050 *Journal of Applied Physiology* **88** 1474–1480.
- 1051 Semenza GL 2012 Hypoxia-inducible factors in physiology and medicine. *Cell* **148** 399–408.
- 1052 Semenza GL 2016 The hypoxic tumor microenvironment: a driving force for breast cancer
 1053 progression. *Biochimica et Biophysica Acta* **1863** 382–391.

- 1054 Sharkey AM, Day K, McPherson A, Malik S, Licence D, Smith SK & Charnock-Jones DS 2000 Vascular
1055 endothelial growth factor expression in human endometrium is regulated by hypoxia. *The Journal of*
1056 *Clinical Endocrinology & Metabolism* **85** 402–409.
- 1057 Shifren JL, Tseng JF, Zaloudek CJ, Ryan IP, Meng YG, Ferrara N, Jaffe RB & Taylor RN 1996 Ovarian
1058 steroid regulation of vascular endothelial growth factor in the human endometrium: implications for
1059 angiogenesis during the menstrual cycle and in the pathogenesis of endometriosis. *The Journal of*
1060 *Clinical Endocrinology & Metabolism* **81** 3112–3118.
- 1061 Short RV 1976 The evolution of human reproduction. *Proceedings of the Royal Society of London.*
1062 *Series B. Biological Sciences* **195** 3–24.
- 1063 Sourbron S 2010 Technical aspects of MR perfusion. *European Journal of Radiology* **76** 304–313.
- 1064 Sourbron SP & Buckley DL 2012 Tracer kinetic modelling in MRI: estimating perfusion and capillary
1065 permeability. *Physics in Medicine and Biology* **57** R1–R33.
- 1066 Stavri GT, Zachary IC, Baskerville PA, Martin JF & Erusalimsky JD 1995 Basic fibroblast growth factor
1067 upregulates the expression of vascular endothelial growth factor in vascular smooth muscle cells.
1068 *Circulation* **92** 11–14.
- 1069 Stephen Charnock-Jones D, Sharkey AM, Rajput-Williams J, Burch D, Paul Schofield J, Fountain SA,
1070 Boocock CA & Smith SK 1993 Identification and localization of alternately spliced mRNAs for vascular
1071 endothelial growth factor in human uterus and estrogen regulation in endometrial carcinoma cell
1072 lines. *Biology of Reproduction* **48** 1120–1128.
- 1073 Stewart E, Cookson C, Gandolfo R & Schulze-Rath R 2017 Epidemiology of uterine fibroids: a
1074 systematic review. *BJOG: An International Journal of Obstetrics & Gynaecology* **124** 1501–1512.
- 1075 Stewart EA, Laughlin-Tommaso SK, Catherino WH, Lalitkumar S, Gupta D & Vollenhoven B 2016
1076 Uterine fibroids. *Nature Reviews Disease Primers* **2** 16043.
- 1077 Suda T, Takubo K & Semenza GL 2011 Metabolic regulation of hematopoietic stem cells in the
1078 hypoxic niche. *Cell Stem Cell* **9** 298–310.
- 1079 Sugino N *et al.* 1996 Changes in activity of superoxide dismutase in the human endometrium
1080 throughout the menstrual cycle and in early pregnancy. *Human Reproduction* **11** 1073–1078.
- 1081 Takahashi N *et al.* 2016 The assessment of myometrium perfusion in patients with uterine fibroid by
1082 arterial spin labeling MRI. *SpringerPlus* **5** 1907.
- 1083 Takeda N *et al.* 2010 Differential activation and antagonistic function of HIF- isoforms in

- macrophages are essential for NO homeostasis. *Genes & Development* **24** 491–501.
- Taylor CT 2008 Interdependent roles for hypoxia inducible factor and nuclear factor- κ B in hypoxic inflammation. *The Journal of Physiology* **586** 4055–4059.
- Taylor CT, Doherty G, Fallon PG & Cummins EP 2016 Hypoxia-dependent regulation of inflammatory pathways in immune cells. *Journal of Clinical Investigation* **126** 3716–3724.
- Thiruchelvam U, Dransfield I, Saunders PTK & Critchley HOD 2013 The importance of the macrophage within the human endometrium. *Journal of Leukocyte Biology* **93** 217–225.
- Thiruchelvam U, Maybin JA, Armstrong GM, Greaves E, Saunders PTK & Critchley HOD 2016 Cortisol regulates the paracrine action of macrophages by inducing vasoactive gene expression in endometrial cells. *Journal of Leukocyte Biology* **99** 1165–1171.
- Thompson AAR *et al.* 2014 Hypoxia-inducible factor 2 α regulates key neutrophil functions in humans, mice, and zebrafish. *Blood* **123** 366–376.
- Velnar T, Bailey T & Smrkolj V 2009 The wound healing process: an overview of the cellular and molecular mechanisms. *Journal of International Medical Research* **37** 1528–1542.
- Walmsley SR *et al.* 2005 Hypoxia-induced neutrophil survival is mediated by HIF-1 α -dependent NF- κ B activity. *The Journal of Experimental Medicine* **201** 105–115.
- Wang GL, Jiang BH, Rue EA & Semenza GL 1995 Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proceedings of the National Academy of Sciences* **92** 5510–5514.
- Wang Q, Xu X, He B, Li Y, Chen X & Wang J 2013 A critical period of progesterone withdrawal precedes endometrial breakdown and shedding in mouse menstrual-like model. *Human Reproduction* **28** 1670–1678.
- Wheeler KC, Jena MK, Pradhan BS, Nayak N, Das S, Hsu C-D, Wheeler DS, Chen K & Nayak NR 2018 VEGF may contribute to macrophage recruitment and M2 polarization in the decidua. *PLOS ONE*. Edited by X. Ye **13** e0191040.
- Wu B *et al.* 2014 ROS are critical for endometrial breakdown via NF- κ B–COX-2 signaling in a female mouse menstrual-like model. *Endocrinology* **155** 3638–3648.
- Wu M-H, Chen K-F, Lin S-C, Lgu C-W & Tsai S-J 2007 Aberrant expression of leptin in human endometriotic stromal cells is induced by elevated levels of hypoxia inducible factor-1 α . *The American Journal of Pathology* **170** 590–598.

Young VJ, Brown JK, Maybin J, Saunders PTK, Duncan WC & Horne AW 2014 Transforming growth factor- β induced warburg-like metabolic reprogramming may underpin the development of peritoneal endometriosis. *The Journal of Clinical Endocrinology & Metabolism* **99** 3450–3459.

Zagzag D, Lukyanov Y, Lan L, Ali MA, Esencay M, Mendez O, Yee H, Voura EB & Newcomb EW 2006 Hypoxia-inducible factor 1 and VEGF upregulate CXCR4 in glioblastoma: implications for angiogenesis and glioma cell invasion. *Laboratory Investigation* **86** 1221–1232.

Zhang J & Salamonsen LA 2002 Expression of hypoxia-inducible factors in human endometrium and suppression of matrix metalloproteinases under hypoxic conditions do not support a major role for hypoxia in regulating tissue breakdown at menstruation. *Human Reproduction* **17** 265–274.

Zhang P, Yao Q, Lu L, Li Y, Chen P-J & Duan C 2014 Hypoxia-inducible factor 3 is an oxygen-dependent transcription activator and regulates a distinct transcriptional response to hypoxia. *Cell Reports* **6** 1110–1121.

Zhang X *et al.* 2007 Wild fulvous fruit bats (*Rousettus leschenaulti*) exhibit human-like menstrual cycle. *Biology of Reproduction* **77** 358–364.

Zheng L, Kelly CJ & Colgan SP 2015 Physiologic hypoxia and oxygen homeostasis in the healthy intestine. A review in the theme: cellular responses to hypoxia. *American Journal of Physiology. Cell Physiology* **309** C350–C360.

FIGURE LEGENDS

Figure 1. Rodent models of simulated menstruation. (a) An exogenous hormone mouse model of simulated menstruation. Female mice are ovariectomized (ovex) and allowed to recover for 7-14 days before being given subcutaneous injections of oestradiol (E_2). A progesterone (P_4) implant is subcutaneously inserted and lower dose E_2 injections administered. The decidualisation stimulus (oil) is intracervically administered. In order to induce a *menstrual-like* event, the P_4 implant is subsequently removed (T_0). This triggers a menstrual like bleed (8h after P_4 withdrawal- T_8) and subsequent endometrial regeneration (24h after P_4 withdrawal- T_{24}). **(b) A pseudopregnancy mouse model of simulated menstruation.** Female mice are mated with vasectomized males to induce pseudopregnancy. Three to four days after the formation of the vaginal plug, decidualisation is externally induced via uterine oil injection. Two days after the decidualisation stimulus, mice are ovariectomized (ovex) to trigger P_4 withdrawal (T_0). Using this approach, endometrial breakdown is apparent

12-16h after P₄ withdrawal (T₁₂) and re-epithelialisation can be detected 24h after P₄ withdrawal. Optionally, mice can receive daily subcutaneous injections of E₂ to prevent atrophy of the uterus following ovariectomy.

(c) Xenograft mouse model. Female immunodeficient mice are ovariectomized (ovex) and allowed to recover for 7-14 days before the implantation of the endometrial patches/dissociated endometrial cells. At the time of implantation or shortly after the formation of the xenografts, mice are treated with E₂ and P₄ for 21-28 days to induce the menstrual cycle. When the P₄ pellet is removed, menstruation and successive regeneration takes place in the xenograft for the next 4-8 days.

Figure 2. In vivo methods with the potential to detect markers of endometrial hypoxia in women. Left: previous *in vivo* work to assess human endometrial hypoxia, shown with structural MRI of the uterus and surrounding tissues. Right: potential non-invasive imaging methods for translation from other body areas. [+] indicates advantages of each technique, [-] indicates disadvantages. Relevant references shown for each. DCE-MRI = Dynamic contrast-enhanced MRI, MRI = Magnetic resonance imaging.

Figure 3. Overview of the presence and role of hypoxia in endometrial physiology. (a) Hypoxia during implantation. Endometrial stromal cells undergo decidualisation under the influence of progesterone. Hypoxia inducible factor (HIF)-2 α in these uterine stromal cells supports decidualisation, embryo invasion and survival. Endometrial blood vessels undergo dynamic remodelling that may be influenced by hypoxia/HIF. **(b) Hypoxia during endometrial breakdown.** Vasoconstriction of the endometrial vessels may limit blood loss during menstruation and cause transient endometrial hypoxia to stabilise HIF-1 α . The endometrial leukocyte population may be altered in number and/or function by hypoxia/HIF. **(c) Role of hypoxia during endometrial repair.** Hypoxia is not detected in endometrial areas that have reepithelialised, while those areas undergoing active regeneration remain hypoxic. This hypoxia is thought to promote endometrial VEGF (alongside other factors), which is responsible for reepithelialisation and new blood vessel formation. P₄ = progesterone, G = glands, BV = blood vessel, VEGF = vascular endothelial growth factor, HIF = hypoxia-inducible factor.

Figure 4. Abnormal uterine bleeding (AUB) and the potential role of hypoxia. Abnormal uterine bleeding may be due to structural (Polyps, Adenomyosis, Leiomyoma, Malignancy) or non-structural (Coagulopathy, Ovulatory, Endometrial, Iatrogenic or Not otherwise classified) disorders. The role of hypoxia in AUB is

unknown but its potential role in four disorders is illustrated. **(I)** Leiomyoma (fibroids): the decreased levels of endothelin and PG2F α receptors may compromise endometrial vasoconstriction and increase menstrual blood flow. **(II)** Malignancy: tumour hypoxia leads to disorganised angiogenesis and increased metastasis. **(III)** Endometrial disorders: endothelial cell focal discontinuities and impairment of vascular smooth muscle cells may influence vasoconstriction. This may decrease HIF-1 α and prevent optimal post-menstrual repair. **(IV)** Adenomyosis: VEGF and HIF-1 α overexpression may contribute to increased vessel formation and AUB. G = glands, BV = blood vessels, Myo = myometrium, E = endometrium, VEGF = vascular endothelial growth factor, HIF = hypoxia-inducible factor.




Figure 1. Rodent models of simulated menstruation. (a) An exogenous hormone mouse model of simulated menstruation. Female mice are ovariectomized (ovex) and allowed to recover for 7-14 days before being given subcutaneous injections of oestradiol (E₂). A progesterone (P₄) implant is subcutaneously inserted and lower dose E₂ injections administered. The decidualisation stimulus (oil) is intracervically administered. In order to induce a menstrual-like event, the P₄ implant is subsequently removed (T₀). This triggers a menstrual like bleed (8h after P₄ withdrawal-T₈) and subsequent endometrial regeneration (24h after P₄ withdrawal-T₂₄). (b) A pseudopregnancy mouse model of simulated menstruation. Female mice are mated with vasectomized males to induce pseudopregnancy. Three to four days after the formation of the vaginal plug, decidualisation is externally induced via uterine oil injection. Two days after the decidualisation stimulus, mice are ovariectomized (ovex) to trigger P₄ withdrawal (T₀). Using this approach, endometrial breakdown is apparent 12-16h after P₄ withdrawal (T₁₂) and re-epithelialisation can be detected 24h after P₄ withdrawal. Optionally, mice can receive daily subcutaneous injections of E₂ to prevent atrophy of the uterus following ovariectomy. (c) Xenograft mouse model. Female immunodeficient mice are ovariectomized (ovex) and allowed to recover for 7-14 days before the implantation of the endometrial patches/dissociated endometrial cells. At the time of implantation or shortly after the formation of the xenografts, mice are treated with E₂ and P₄ for 21-28 days to induce the menstrual cycle. When the P₄ pellet is removed, menstruation and successive regeneration takes place in the xenograft for the next 4-8 days.

178x156mm (150 x 150 DPI)




Figure 2. In vivo methods with the potential to detect markers of endometrial hypoxia in women. Left: previous in vivo work to assess human endometrial hypoxia, shown with structural MRI of the uterus and surrounding tissues. Right: potential non-invasive imaging methods for translation from other body areas. [+] indicates advantages of each technique, [-] indicates disadvantages. Relevant references shown for each. DCE-MRI = Dynamic contrast-enhanced MRI, MRI = Magnetic resonance imaging.

338x190mm (225 x 225 DPI)




Figure 3. Overview of the presence and role of hypoxia in endometrial physiology. (a) Hypoxia during implantation. Endometrial stromal cells undergo decidualisation under the influence of progesterone. Hypoxia inducible factor (HIF)-2 α in these uterine stromal cells supports decidualisation, embryo invasion and survival. Endometrial blood vessels undergo dynamic remodelling that may be influenced by hypoxia/HIF. (b) Hypoxia during endometrial breakdown. Vasoconstriction of the endometrial vessels may limit blood loss during menstruation and cause transient endometrial hypoxia to stabilise HIF-1 α . The endometrial leukocyte population may be altered in number and/or function by hypoxia/HIF. (c) Role of hypoxia during endometrial repair. Hypoxia is not detected in endometrial areas that have reepithelialised, while those areas undergoing active regeneration remain hypoxic. This hypoxia is thought to promote endometrial VEGF (alongside other factors), which is responsible for reepithelialisation and new blood vessel formation. P4 = progesterone, G = glands, BV = blood vessel, VEGF = vascular endothelial growth factor, HIF = hypoxia-inducible factor.




Figure 4. Abnormal uterine bleeding (AUB) and the potential role of hypoxia. Abnormal uterine bleeding may be due to structural (Polyps, Adenomyosis, Leiomyoma, Malignancy) or non-structural (Coagulopathy, Ovulatory, Endometrial, Iatrogenic or Not otherwise classified) disorders. The role of hypoxia in AUB is unknown but its potential role in four disorders is illustrated. (I) Leiomyoma (fibroids): the decreased levels of endothelin and PG2Fa receptors may compromise endometrial vasoconstriction and increase menstrual blood flow. (II) Malignancy: tumour hypoxia leads to disorganised angiogenesis and increased metastasis. (III) Endometrial disorders: endothelial cell focal discontinuities and impairment of vascular smooth muscle cells may influence vasoconstriction. This may decrease HIF-1α and prevent optimal post-menstrual repair. (IV) Adenomyosis: VEGF and HIF-1α overexpression may contribute to increased vessel formation and AUB. G = glands, BV = blood vessels, Myo = myometrium, E = endometrium, VEGF = vascular endothelial growth factor, HIF = hypoxia-inducible factor.